

Closures of Conjugacy Classes in G_2

HANSPETER KRAFT*

*Mathematisches Institut, Universität Basel, Rheinsprung 21,
CH-4051 Basel, Switzerland*

Communicated by Claudio Procesi

Received June 10, 1988

1. INTRODUCTION

1.1. In a recent paper [LS] Levasseur and Smith have shown that the 8-dimensional nilpotent conjugacy class in the simple Lie algebra \mathfrak{g} of type G_2 has a non-normal closure. In the following we give a short proof of this result and show that all other classes have a normal closure. This completes the geometric picture in the spirit of the work [KP1, KP2, KP3], which deals with the case of the classical groups.

1.2. THEOREM.¹ *Let \mathfrak{g} be the simple Lie algebra of type G_2 and let C_i be the nilpotent conjugacy class in \mathfrak{g} of dimension $i = 6, 8, 10$, and 12.*

(a) *Every conjugacy class of \mathfrak{g} except C_8 has a normal closure with rational singularities.*

(b) *[Levasseur and Smith] $\overline{C_8}$ is not normal in $\overline{C_6} = \overline{C_8} \setminus C_8$. The normalization $\eta_8: \widetilde{C_8} \rightarrow \overline{C_8}$ is bijective and $\widetilde{C_8}$ has an isolated rational singularity in $\eta_8^{-1}(0)$.*

(c) *$\overline{C_{12}}$ has a singularity of type D_4 in C_{10} , and $\overline{C_{10}}$ a singularity of type A_1 in C_8 .*

The proof of these results is based on the same construction as in [LS]: We embed \mathfrak{g} into \mathfrak{so}_7 by the 7-dimensional standard representation, and study the \mathfrak{g} -equivariant projection $p: \mathfrak{so}_7 \rightarrow \mathfrak{g}$. It turns out that the restriction of p to the 8- and 10-dimensional nilpotent conjugacy classes D_8 and D_{10} in \mathfrak{so}_7 induces finite surjective morphisms

$$p_8: \overline{D_8} \rightarrow \overline{C_8} \quad \text{and} \quad p_{10}: \overline{D_{10}} \rightarrow \overline{C_{10}}.$$

The result then follows from a careful analysis of these two maps.

* Partially supported by Schweizerischer Nationalfonds.

¹ In a preliminary version of this paper under the title "Non-normality of Closures of Conjugacy Classes in G_2 " the statement of this theorem is not correct.

1.3. *Remark.* The simple group G_2 also has an exceptional behavior with respect to the *sheets* in its Lie algebra \mathfrak{g} (cf. [Kr1], [BK], or [B] for definitions). One of the two subregular sheets through C_{10} is smooth, the other is not normal and C_{10} undergoes a threefold covering in the normalization of that sheet [Sl, p. 151; BK, 7.3 Beispiel b].

Conventions. The base field k is algebraically closed of characteristic zero. If X is an algebraic variety we denote by $k[X]$ the algebra of global regular functions on X .

2. NILPOTENT CONJUGACY CLASSES IN G_2

2.1. Let G be a simple group of type G_2 . Fix a maximal torus T and a Borel subgroup $B \supset T$ and denote by α_1, α_2 the corresponding base of the root system Φ with respect to T , where α_1 is a short and α_2 a long root. The nilpotent cone of the Lie algebra $\mathfrak{g} := \text{Lie } G$ consists of five conjugacy classes C_{12}, C_{10}, C_8, C_6 , and $C_0 = \{0\}$ with dimensions $\dim C_i = i$. C_6 is the conjugacy class of a long root vector $x_2 \in \mathfrak{g}_{\alpha_2}$, C_8 the class of a short root vector $x_1 \in \mathfrak{g}_{\alpha_1}$, C_{10} the class of $x_2 + x'_2$, where $x'_2 \in \mathfrak{g}_{3\alpha_1 + \alpha_2}$ is another long root vector, and C_{12} the class of $x_1 + x_2$. C_{12} is the *regular class*, C_{10} the *subregular class*, and we have $\overline{C_i} \supset C_j$ for $i \geq j$. (Cf. [SK])

2.2. The long root vectors generate the subalgebra

$$\text{Lie } T \oplus \bigoplus_{\beta \text{ long}} \mathfrak{g}_{\beta}$$

of \mathfrak{g} which we will identify with \mathfrak{sl}_3 . It is easy to see that \mathfrak{g} decomposes as an \mathfrak{sl}_3 -module in the form

$$\mathfrak{g} = \mathfrak{sl}_3 \oplus k^3 \oplus (k^3)^*.$$

We can therefore consider SL_3 as a subgroup of G .

2.3. *Remark.* Let $H := G_x$ be the centralizer of $x := x_2 + x'_2 \in C_{10}$. It is known that H^0 is unipotent and that $H/H^0 \simeq \mathcal{S}_3$, the symmetric group in three letters (see [Ca, table on p. 401]). We will only need that H contains the center $Z \simeq \mathbb{Z}/3\mathbb{Z}$ of SL_3 and that

$$\text{Lie } H \supset \mathfrak{g}_{\alpha_1 + \alpha_2} \oplus \mathfrak{g}_{2\alpha_1 + \alpha_2} \oplus \mathfrak{g}_{3\alpha_1 + 2\alpha_2};$$

this is obvious from what we have said above.

2.4. *Remark.* The centralizer of $x_1 \in C_8$ is connected (see [Ca, table on p. 401]).

3. THE STANDARD REPRESENTATION OF G_2

3.1. Let $\rho: G \rightarrow GL(V)$ be the 7-dimensional irreducible representation with highest weight $\omega_1 = 2\alpha_1 + \alpha_2$. This representation is orthogonal, and the invariant quadratic form $q: V \rightarrow k$ generates the algebra $k[V]^G$ of G -invariant polynomials on V . The zero cone $V^\circ := q^{-1}(0)$ consists of two orbits, $\{0\}$ and Gv_0 , where $v_0 \in V_{\omega_1}$ is a highest weight vector. (As usual we denote by V_β the weight space of V of weight β .)

The representation ρ defines an embedding $G \hookrightarrow SO_7$. The adjoint representation of G on \mathfrak{so}_7 is isomorphic to $\wedge^2 V$ and decomposes in the form

$$\mathfrak{so}_7 \simeq \bigwedge^2 V \simeq \mathfrak{g} \oplus V.$$

(This is clear from dimensional reason.)

The weights of V are the zero weight and the short roots of \mathfrak{g} , all with multiplicity one. Hence we have

$$V \simeq k \oplus k^3 \oplus (k^3)^*$$

as an SL_3 -module.

3.2. LEMMA. *Let $v_0 \in V \subset \mathfrak{so}_7$ be a highest weight vector of V . As an element of \mathfrak{so}_7 the endomorphism v_0 is nilpotent with partition $(3, 2, 2)$.*

(The *partition* of a nilpotent endomorphism is given by the sizes of the blocks in a Jordan normal form.)

Proof. The highest weight of V is $\omega_1 := 2\alpha_1 + \alpha_2$, and the corresponding weight space in $\wedge^2 V$ is 2-dimensional:

$$\left(\bigwedge^2 V \right)_{\omega_1} = V_0 \wedge V_{\omega_1} \oplus V_{\alpha_1} \wedge V_{\alpha_1 + \alpha_2}.$$

Since a highest weight vector $v_0 \in V \subset \wedge^2 V$ is annihilated by \mathfrak{g}_{α_1} and \mathfrak{g}_{α_2} we see that v_0 has non-zero components in both summands:

$$v_0 = w_0 \wedge w_2 + w_1 \wedge w_3,$$

$$w_0 \in V_0, w_1 \in V_{\alpha_1}, w_2 \in V_{2\alpha_1 + \alpha_2}, w_3 \in V_{\alpha_1 + \alpha_2} \text{ and all } w_i \neq 0.$$

Furthermore, the G -isomorphism $\sigma: V \xrightarrow{\sim} V^*$ satisfies $\sigma(V_\beta) = (V_{-\beta})^*$ for all weights β . It follows that the composition

$$\bigwedge^2 V \hookrightarrow V \otimes V \xrightarrow{\sim} V \otimes V^* \xrightarrow{\sim} \text{End } V$$

maps the element

$$w_0 \wedge w_2 + w_1 \wedge w_3 = \frac{1}{2}(w_0 \otimes w_2 - w_2 \otimes w_0 + w_1 \otimes w_3 - w_3 \otimes w_1)$$

to an element of the form

$$w = w_0 \otimes \bar{w}_{-2} + w_2 \otimes \bar{w}_0 + w_1 \otimes \bar{w}_{-3} + w_3 \otimes \bar{w}_{-1}$$

with non-zero elements $\bar{w}_0 \in (V_0)^*$, $\bar{w}_{-1} \in (V_{-\alpha_1})^*$, $\bar{w}_{-2} \in (V_{-2\alpha_1-\alpha_2})^*$, $\bar{w}_{-3} \in (V_{-\alpha_1-\alpha_2})^*$. It is now easy to see that w is a nilpotent element of $\text{End } V$ with partition $(3, 2, 2)$. ■

4. THE FUNDAMENTAL CONSTRUCTION

4.1. The nilpotent cone of \mathfrak{so}_7 consists of the conjugacy classes D_{18} , D_{16} , D_{14} , D_{12} , D_{10} , D_8 , and $D_0 = \{0\}$ of dimensions $\dim D_i = i$. They are completely determined by their conjugacy class in $M_7(k)$ with respect to $GL_7(k)$ and correspond to the partitions (7) , $(5, 1, 1)$, $(3, 3, 1)$, $(3, 2, 2)$, $(3, 1^4)$, $(2, 2, 1^3)$, and (1^7) . (For this and the following see [KP3, Sect. 19 tables].) We have $\overline{D_i} \supset D_j$ for $i \geq j$, and all D_i except D_{12} have a normal closure $\overline{D_i}$ with rational singularities.

Consider the G -linear projection $p: \mathfrak{so}_7 \rightarrow \mathfrak{g}$ given by the decomposition $\mathfrak{so}_7 = \mathfrak{g} \oplus V$ (see Section 3).

4.2. PROPOSITION. *The map p induces finite surjective morphisms*

$$p_8: \overline{D_8} \rightarrow \overline{C_8} \quad \text{and} \quad p_{10}: \overline{D_{10}} \rightarrow \overline{C_{10}}.$$

The morphism p_8 is bijective, but it is not an isomorphism in the points of $p_8^{-1}(\overline{C_6})$ (i.e., the fibres over these points are not reduced).

Proof. By the lemma above we have $\overline{D_{10}} \cap V = \overline{D_8} \cap V = \{0\}$. Since the closures $\overline{D_i}$ are closed G -stable cones in \mathfrak{so}_7 it follows that the images $X_8 := p(\overline{D_8})$ and $X_{10} := p(\overline{D_{10}})$ are closed and G -stable cones in \mathfrak{g} and that the maps $p_8: \overline{D_8} \rightarrow X_8$ and $p_{10}: \overline{D_{10}} \rightarrow X_{10}$ are finite morphisms. In fact, given a finitely generated graded algebra $R = \bigoplus_i R_i$ with $R_0 = k$ and a graded subalgebra $S = \bigoplus_i S_i$ such that $\sqrt{RS_+} = R_+$, where R_+ and S_+ are the homogeneous maximal ideals of R and S , then R is a finitely generated S -module (see, for example, [Kr2, II.4.3 Satz 8] and its proof). But C_6 and C_8 are the only conjugacy classes in \mathfrak{g} of dimension 6 and 8, all other classes have dimension ≥ 10 (see Section 2). Hence $X_8 = \overline{C_8}$, and X_{10} contains a dense 10-dimensional conjugacy class and so $X_{10} = \overline{C_{10}}$. Since the centralizer of $x_1 \in C_8$ is connected (Remark 2.4) the map p_8 is birational.

Let $x_2 \in C_6$ be a long root vector as in Section 2. We claim that $p_8^{-1}(x_2) = \{(x_2, 0)\} \subset \mathfrak{g} \oplus V$. In particular $p_8^{-1}(\overline{C_6}) \rightarrow \overline{C_6}$ is bijective. In fact, consider the B -stable line $kx_2 \subset \mathfrak{g}$. Then $p_8^{-1}(kx_2)$ is finite union of lines, each one stable under B . Since there are only two B -stable lines in \mathfrak{so}_7 , namely $k(x_2, 0)$ and the highest weight space of V , the claim follows.

Next we show that the fibre $p_8^{-1}(x_2)$ is not reduced. In fact, the fibre $p_8^{-1}(x_2)$ is the schematic intersection of D_8 with $\{x_2\} \times V$. For the tangent spaces we find

$$T_{(x_2, 0)} D_8 = [x_2, \mathfrak{so}_7] = [x_2, \mathfrak{g}] \oplus [x_2, V].$$

Since $[x_2, V] \neq (0)$ the intersection $D_8 \cap (x_2, V)$ is not transversal, hence the fibre $p_8^{-1}(x_2)$ is not a reduced point. ■

4.3. *Remark.* Every nilpotent class $C_i \subset \mathfrak{g}$ generates a nilpotent class in \mathfrak{so}_7 , via the embedding $\mathfrak{g} \subset \mathfrak{so}_7$ (3.1). From the explicit description of the nilpotent classes C_i in 2.1 it is easy to determine the nilpotent endomorphism of V induced by $x \in C_i$; e.g., a short root $x_2 \in \mathfrak{g}$ defines an endomorphism with partition $(3, 2, 2)$ and a long root $x_2 \in \mathfrak{g}$ one with partition $(2, 2, 1^3)$. Using 4.1 we find the following inclusions:

$$C_6 \subset D_8, \quad C_8 \subset D_{12}, \quad C_{10} \subset D_{14}, \quad C_{12} \subset D_{18}.$$

5. MULTIPLICITIES

5.1. For any G -variety X we denote by $\text{mult}_M(X)$ the multiplicity of an irreducible representation M in the algebra $k[X]$ of global regular functions on X , i.e.,

$$\text{mult}_M(X) = \dim_k \text{Mor}_G(X, M^*),$$

where $\text{Mor}_G(X, M^*)$ is the k -vectorspace of G -equivariant morphisms $X \rightarrow M^*$ into the dual module M^* of M . If X is a G -orbit, $X \simeq G/H$, we obtain

$$\text{mult}_M(X) = \dim(M^*)^H \quad (\text{Frobenius reciprocity}).$$

Now let $C \subset \mathfrak{g}$ be a conjugacy class and \bar{C} its closure in \mathfrak{g} . Since the complement $\bar{C} \setminus C$ is of codimension ≥ 2 we have the following result due to Kostant ([Ko, 2.2 Proposition 9]; cf. [BK]):

PROPOSITION. *The closure \bar{C} of the conjugacy class C is normal if and only if $\text{mult}_M(\bar{C}) = \text{mult}_M(C)$ for all irreducible representations M of G .*

5.2. For the proof of the normality of $\overline{C_{10}}$ we will need to know the multiplicities of the representations V and W of highest weight ω_1 and $2\omega_1$ in $k[C_{10}]$.

LEMMA. *For the representations V and W of highest weight ω_1 and $2\omega_1$ we have*

$$\text{mult}_V(C_{10}) = 0 \quad \text{and} \quad \text{mult}_W(C_{10}) \leq 1.$$

Proof. We know that C_{10} is the conjugacy class of $x = x_2 + x'_2$ (notations of 2.1). By Frobenius reciprocity (5.1) we have to show that

$$V^H = 0 \quad \text{and} \quad \dim W^H \leq 1,$$

where $H = G_x$ is the centralizer of x . (Remember that V and W are selfdual.)

Let $U \subset G$ be the unipotent subgroup with Lie algebra

$$\text{Lie } U = \mathfrak{g}_{\alpha_1 + \alpha_2} \oplus \mathfrak{g}_{2\alpha_1 + \alpha_2} \oplus \mathfrak{g}_{3\alpha_1 + 2\alpha_2}.$$

We know that $U \subset H$ (Remark 2.3). Since U is normalized by the maximal torus T the fixed point set V^U is a sum of weight spaces. In fact, it is easy to see that

$$V^U = V_{\alpha_1 + \alpha_2} \oplus V_{2\alpha_1 + \alpha_2}.$$

Now the center Z of SL_3 belongs to H , too (Remark 2.3). Since every short root β is non-trivial on Z we obtain

$$V^H \subset (V^U)^Z = 0,$$

proving the first claim.

The second symmetric power $S^2(V)$ of V contains the irreducible representation W , which is of dimension 27, hence $S^2(V) \simeq k \oplus W$ as a G -module. It follows that the weights in W are 0, α_1 , α_2 , $2\alpha_1$ and their conjugates under the Weyl group, and their multiplicities are given by

$$\dim W_0 = 3, \quad \dim W_{\alpha_1} = 2, \quad \dim W_{\alpha_2} = 1, \quad \dim W_{2\alpha_1} = 1.$$

It is easy to see that

$$W^U \subset W_0 \oplus W_{\alpha_1 + \alpha_2} \oplus W_{2\alpha_1 + \alpha_2} \oplus W_{2(\alpha_1 + \alpha_2)} \oplus W_{2(2\alpha_1 + \alpha_2)} \oplus W_{3\alpha_1 + 2\alpha_2}.$$

We claim that $W^U \cap W_0 = 0$, and therefore

$$W^U \subset W_{\alpha_1 + \alpha_2} \oplus \cdots \oplus W_{3\alpha_1 + 2\alpha_2}.$$

In fact, if $w \in W_0$ is annihilated by \mathfrak{g}_β , it is also annihilated by $\mathfrak{g}_{-\beta}$. It follows for an element $w \in W^U \cap W_0$ that $\mathfrak{g}_\beta w = 0$ for every short root β , hence w is annihilated by all of \mathfrak{g} , and the claim follows.

Considering now the action of the center Z of SL_3 the same argument as above applies and shows that

$$W^H \subset (W^U)^Z \subset W_{3\alpha_1 + 2\alpha_2},$$

proving the second claim. ■

6. NORMALITY OF $\overline{C_{10}}$

6.1. We want to calculate the schematic fibre of $0 \in \overline{C_{10}}$ under the finite surjective morphism $p_{10}: \overline{D_{10}} \rightarrow \overline{C_{10}}$ (4.2). By construction, $p_{10}^{-1}(0)$ is the schematic intersection $\overline{D_{10}} \cap V$, hence of the form $\text{Spec } k[V]/J$, where J is a graded G -stable ideal whose radical is the homogeneous maximal ideal of $k[V]$.

6.2. LEMMA. *The ideal J contains the invariant q and all homogeneous elements of degree ≥ 3 . In particular, the only representations possibly occurring in $k[V]/J$ are k , V , and W .*

Proof. Clearly $q \in J$ since q is the restriction of the non-degenerate quadratic invariant $\tilde{q} \in k[\mathfrak{so}_7]^{\text{SO}_7}$ to $V \subset \mathfrak{so}_7$, and \tilde{q} vanishes on D_{10} . Furthermore, the elements of $\overline{D_{10}} \subset \mathfrak{so}_7$ are endomorphisms of rank ≤ 2 (see 4.1). Hence the 3×3 -minors generate a \mathfrak{so}_7 -stable subvector space $M \subset k[\mathfrak{so}_7]$ of homogeneous polynomials of degree 3 vanishing on $\overline{D_{10}}$. More precisely $\overline{D_{10}}$ is the zero set of M together with all homogeneous SO_7 -invariant functions on \mathfrak{so}_7 . These invariants restrict to multiples of q on V (3.1). Since $\overline{D_{10}} \cap V = \{0\}$ this implies that q and the restriction $\bar{M} := M|_V$ generate an ideal of finite index in $k[V]$.

Now the zero fibre $q^{-1}(0) \subset V$ is the closure of the orbit of a highest weight vector, hence $k[q^{-1}(0)] \simeq \bigoplus_{i \geq 0} V_i$ as a G -module, where V_i is simple of highest weight $i\omega_1$ [VP, Theorem 2]. It follows that

$$k[V] \simeq k[q] \otimes \bigoplus_{i \geq 0} V_i$$

as a G -module because q is an irreducible polynomial and V a free $k[q]$ -module (cf. [Ko]). In particular,

$$k[V]_0 = k, \quad k[V]_1 \simeq V, \quad k[V]_2 \simeq kq \oplus W, \quad k[V]_3 \simeq qV \oplus V_3,$$

where $k[V]_d$ is the homogeneous component of $k[V]$ of degree d . Now $\bar{M} \subseteq k[V]_3$ is G -stable and $\bar{M} \not\subseteq qV$, since q and \bar{M} generate an ideal of finite index, as we have seen above. Hence $\bar{M} \supseteq V_3$ and so

$$J \supseteq (q, \bar{M}) \supseteq \bigoplus_{i \geq 3} k[V]_i. \quad \blacksquare$$

6.3. Proof of Normality. Now we want to show that $\overline{C_{10}}$ is normal. Let $\eta: \widetilde{C_{10}} \rightarrow \overline{C_{10}}$ be the normalization. Since $\overline{D_{10}}$ is normal [KP3, Sect. 19 tables], we get a factorization

$$p_{10}: \overline{D_{10}} \xrightarrow{\widetilde{p}_{10}} \widetilde{C_{10}} \xrightarrow{\eta} \overline{C_{10}}$$

with a finite surjective morphism \widetilde{p}_{10} . In terms of coordinate rings this means that we have finite extensions

$$k[\overline{C_{10}}] \subset k[\widetilde{C_{10}}] \subset k[\overline{D_{10}}].$$

Next we show that $k[\widetilde{C_{10}}]$ is a direct summand of $k[\overline{D_{10}}]$. By Frobenius reciprocity (5.1), the second inclusion is of the form $k[G]^{H_2} \subset k[G]^{H_1}$ with subgroups $H_1 \subset H_2 \subset G$, and $k[G]^{H_1} \subset k[G]^{H_1^0} = k[G]^{H_2^0}$. Now there is a finite subgroup $F \subset H_2$ such that $H_2 = F \cdot H_2^0$, hence $k[G]^{H_2} = (k[G]^{H_2^0})^F$. This implies that $k[G]^{H_2}$ is a direct summand of $k[G]^{H_2^0}$, and therefore $k[\widetilde{C_{10}}]$ is a direct summand of $k[\overline{D_{10}}]$.

As a consequence of this we see that the schematic fibre $\eta^{-1}(0) \simeq \text{Spec } R$ is given by a G -stable subalgebra $R \subset k[V]/J$ (notations of 6.1). We have to show that $R = k$. But $\text{mult}_\nu(\widetilde{C_{10}}) = \text{mult}_\nu(C_{10}) = 0$ by Lemma 5.2 and so V cannot occur in R . Also $\text{mult}_w(\widetilde{C_{10}}) = \text{mult}_w(C_{10}) \leq 1$ (Lemma 5.2) and $\text{mult}_w(\overline{C_{10}}) \geq 1$ by the lemma below. This shows that R does not contain W either, proving our claim. \blacksquare

6.4. LEMMA. *The map $\mathfrak{so}_7 \rightarrow M_7(k)$, $X \mapsto X^2$, induces a non-trivial G -equivariant morphism $\overline{C_{10}} \rightarrow V$. In particular $\text{mult}_\nu(\overline{C_{10}}) \geq 1$.*

Proof. It follows from Lemma 3.2 that the map $\mathfrak{so}_7 \xrightarrow{\eta^2} M_7(k)$ is non-zero on C_{10} . Its image lies in Sym_7 , the symmetric matrices in $M_7(k)$, and

$$\text{Sym}_7 \simeq S^2 V \simeq V \oplus k$$

as a G -module. If we compose the G -equivariant map

$$\lambda: \mathfrak{g} \hookrightarrow \mathfrak{so}_7 \xrightarrow{\eta^2} \text{Sym}_7$$

with the projection onto k we obtain a quadratic invariant, hence a multiple of q_2 , which vanishes on all nilpotents. It follows that the composition of λ with the other projection, the one onto V , induces a non-trivial covariant $\overline{C}_{10} \rightarrow V$. ■

7. END OF PROOF OF THEOREM 1.2

We first remark that all non-nilpotent conjugacy classes have a normal closure with rational singularities: Every such closure is a G -fibre bundle over a semisimple class, whose fibre is the closure of a nilpotent class in some strict Levi subalgebra of \mathfrak{g} (see [Sl, Lemma 3.10]). The regular class C_{12} has a normal closure with rational singularities [Ko, He1]) and the singularity of \overline{C}_{12} in C_{10} is of type D_4 (see [Sl]). The minimal class C_6 has a normal closure $\overline{C}_6 = C_6 \cup \{0\}$ with rational singularities [Ke].

In the preceding section we have already shown that \overline{C}_{10} is normal. In addition, we have seen in Section 4 that $C_{10} \subset D_{14}$ and that $C_8 \subset D_{12}$ (Remark 4.3). Since \overline{C}_{10} is normal and since the codimensions of C_8 in \overline{C}_{10} and of D_{12} in \overline{D}_{14} are equal we can apply [KP3, Corollary 13.3]. This result states that \overline{C}_{10} has in C_8 a singularity of the same type as \overline{D}_{14} in D_{12} , which is a singularity of type A_1 [KP3, Sect. 19 tables].

Now \overline{D}_{10} is normal with rational singularities (4.1). Applying Boutot's theorem [Bo] to the finite morphism $p_{10}: \overline{D}_{10} \rightarrow \overline{C}_{10}$ it follows that \overline{C}_{10} has rational singularities, too. (In fact we have seen in 6.3 that the coordinate ring $k[\overline{C}_{10}]$ is a direct summand of $k[\overline{D}_{10}]$.)

Finally, $\overline{D}_8 = D_8 \cup \{0\}$ is normal with an isolated rational singularity in 0 (4.1). Therefore the bijective morphism $p_8: \overline{D}_8 \rightarrow \overline{C}_8$ (Proposition 4.2) is the normalization. This finishes the proof of the theorem.

8. SURVEY OF RESULTS ON THE NORMALITY PROBLEM

In this last section we give a brief summary of what is known about normality of closures of conjugacy classes in reductive groups.

8.1. Reduction to the Simple Case. Let G be a reductive group and $\mathfrak{g} = \text{Lie } G$ its Lie algebra. A conjugacy class C in G (or in \mathfrak{g}) is of the form $G *^H C'$, where $H \subseteq G$ is a Levi subgroup and $C' \subseteq \text{Lie } H$ a nilpotent conjugacy class, and the closure \overline{C} of C is isomorphic to $G *^H \overline{C}'$ (cf. [Sl, II.3.10] or [KP3, 0.2]). Furthermore, a class $C \subset \mathfrak{g}_1 \oplus \mathfrak{g}_2$ is a product $C_1 \times C_2$ of two classes, and its closure is $\overline{C}_1 \times \overline{C}_2$. This reduces the problem to the study of nilpotent classes in simple Lie algebras.

Recall that a semisimple group contains only finitely many nilpotent conjugacy classes, and every conjugacy class has even dimension (cf. [BC]).

8.2. Some General Normality Results. Kostant showed in his fundamental paper [Ko] that the *nilpotent cone* in \mathfrak{g} , which is the closure of the *regular nilpotent class* C_{reg} , is a normal complete intersection, and Hesselink proved that it has rational singularities [He1].

For the *minimal class* C_{min} , i.e., the orbit of highest weight vectors in \mathfrak{g} , a general result of Vinberg and Popov [VP] implies that the closures $\bar{C}_{\text{min}} = C_{\text{min}} \cup \{0\}$ are normal, and it follows from Kempf [Ke] that it has rational singularities.

Using Serre's normality criterion Hesselink showed that certain classes in SL_n can be obtained as normal complete intersections in determinantal subvarieties of \mathfrak{g} defined by rank conditions [He2, 1.2]. Hence they are *normal and Cohen-Macaulay*. In addition, using resolution of singularities and [Ke] he discovered several "small" nilpotent classes in \mathfrak{sl}_n , \mathfrak{so}_n and \mathfrak{sp}_n , and in the exceptional Lie algebras besides the regular and the minimal class which have a normal closure with rational singularities, e.g., one in F_4 , one in E_6 , two in E_7 , and one in E_8 .

Brieskorn studied the singularity of the nilpotent cone in the *subregular class* (this is the only class of codimension 2) in case of a simple Lie algebra of type A , D , and E and showed that it is equivalent to a simple surface singularity of corresponding type (cf. Slodowy's book [Sl], where this is extended to all simple Lie algebras).

8.3. Special Linear Groups. In these groups the closure of every conjugacy class is normal and has rational singularities [KP1]. There is a simple algorithm in terms of the partition associated to a nilpotent class C to determine the classes occurring in the closure \bar{C} . Also one can read from the partition the type of the singularity of \bar{C} in the open classes in the boundary $\partial C := \bar{C} \setminus C$, the so-called *minimal singularities* of C [KP2]. This generalizes the results of Brieskorn and Slodowy (8.2).

8.4. Orthogonal and Symplectic Groups. For these groups and their Lie algebras there exist conjugacy classes with non-normal closures [KP3]. Again the partition of a nilpotent conjugacy class C determines which classes appear in the closure \bar{C} , the type of the minimal singularities, and whether \bar{C} is normal or not, except for the so-called *very even* classes. (These are the conjugacy classes in \mathfrak{so}_{4n} which are not stable under the full orthogonal group O_{4n} .) There are partial results about Cohen-Macaulayness and rational singularities.

It is interesting to remark at this point that the non-normal closures \bar{C} are always *branched in codimension 2*, i.e., there is a class $D \subset \bar{C}$ of

codimension 2 which undergoes a 2-fold covering in the normalization $\eta: \tilde{C} \rightarrow \bar{C}$. As we have seen in 1.2 this is not the case for the class C_8 in G_2 .

8.5. *Branched Non-normality (Beynon and Spaltenstein).* In the paper [BS, Sect. 5(E)] one finds the following result (based on a remark of Verdier and Brylinski):

PROPOSITION. *Let $x, y \in \mathfrak{g}$ be nilpotent elements and let \mathcal{B}_y be the fibre of y under the Springer resolution of singularities of the nilpotent cone of \mathfrak{g} . Denote by ρ_x the Weyl-group representation corresponding to x (and the trivial character of the component group G_x/G_x^0) under the Springer-correspondence. Then*

$$\text{mult}_{\rho_x} H^{2\beta(x)}(\mathcal{B}_y) = \# \eta^{-1}(y),$$

where $\beta(x) = \dim \mathcal{B}_x$ and $\eta: \tilde{C} \rightarrow \bar{C}$ is the normalisation.

It follows that these multiplicities determine the inclusion behavior of closures of nilpotent conjugacy classes in \mathfrak{g} and allow one to detect all classes with *branched non-normal closures*. The multiplicities have been calculated by Shoji [Sh] for F_4 and by Beynon and Spaltenstein [BS] for E_6, E_7 , and E_8 . Inspecting their tables one finds the following classes with a branched non-normal closure (we use the notations of Bala and Carter [BC]):

$$F_4: A_1 + B_2, C_3$$

$$E_6: A_4, 2A_2, A_2 + A_1$$

$$E_7: D_6(a_1), A_5'', A_4, A_3 + A_2, D_4(a_1) + A_1$$

$$E_8: E_7(a_1), E_6, E_6(a_1), E_7(a_4), A_6, D_6(a_1), D_5 + A_1, E_7(a_5), A_4, \\ A_3 + A_2, D_4, D_4(a_1), A_3 + A_1, 2A_2 + A_1$$

8.6. *The Method of Richardson.* The paper [Ri] contains a method to calculate the rank of the quotient map $\pi: \mathfrak{g} \rightarrow k^r$ in a given nilpotent element $x \in \mathfrak{g}$. It is easy to see that $\text{rank } d\pi_x = \text{mult}_{\mathfrak{g}}(\overline{C_x})$ (notations 5.1), where C_x is the conjugacy class of x [Ri, Proposition 2.7]. On the other hand, it is shown in [BK] that under certain assumptions about the stabilizer G_x , the multiplicities are constant along the sheets containing the class C_x in case $\overline{C_x}$ is normal. Thus comparing $\text{rank } d\pi_x$ with the rank of the quotient map in the corresponding element of the Levi subalgebra associated to the sheet, Richardson obtains a sufficient criterion for non-normality. He detects in this way

$$4 \text{ classes in } E_6, 5 \text{ classes in } E_7 \text{ and } 11 \text{ classes in } E_8,$$

which have a non-normal closure with a *bijective normalisation*.

ACKNOWLEDGMENTS

I thank Friedrich Knop and Nicola Spaltenstein for helpful discussions.

REFERENCES

- [BC] P. BALA AND R. W. CARTER, Classes of unipotent elements in simple algebraic groups. *Math. Proc. Cambridge Philos. Soc.* **79** (1976), 401–425; **80** (1976), 1–18.
- [BS] W. M. BEYNON AND N. SPALTENSTEIN, Green functions of finite Chevalley groups of type E_n ($n = 6, 7, 8$), *J. Algebra* **88** (1984), 584–614.
- [B] W. BORHO, Über Schichten halbeinfacher Lie-Algebren, *Invent. Math.* **65** (1981), 283–317.
- [BK] W. BORHO AND H. KRAFT, Über Bahnen und deren Deformationen bei linearen Aktionen reduktiver Gruppen, *Comment. Math. Helv.* **54** (1979), 61–104.
- [Bo] J.-F. BOUTOT, Singularités rationnelles et quotients par les groupes réductifs, *Invent. Math.* **88** (1987), 65–68.
- [Ca] R. W. CARTER, “Finite Groups of Lie Type: Conjugacy Classes and Complex Characters,” Wiley–Interscience, Chichester/New York, 1985.
- [He1] W. HESSELINK, Cohomology and the resolution of the nilpotent variety, *Math. Ann.* **223** (1976), 249–251.
- [He2] W. HESSELINK, The normality of closures of orbits in a Lie algebra, *Comment. Math. Helv.* **54** (1979), 105–110.
- [Ke] G. KEMPF, On the collapsing of homogeneous bundles, *Invent. Math.* **37** (1979), 229–239.
- [Ko] B. KOSTANT, Lie group representation on polynomial rings, *Amer. J. Math.* **85** (1963), 327–404.
- [Kr1] H. KRAFT, Parametrisierung von Konjugationsklassen in \mathfrak{sl}_n , *Math. Ann.* **234** (1978), 209–220.
- [Kr2] H. KRAFT, “Geometrische Methoden in der Invariantentheorie,” Vieweg, Braunschweig, 1985.
- [KP1] H. KRAFT AND C. PROCESI, Closures of conjugacy classes of matrices are normal, *Invent. Math.* **53** (1979), 227–247.
- [KP2] H. KRAFT AND C. PROCESI, Minimal singularities in GL_n , *Invent. Math.* **62** (1981), 503–515.
- [KP3] H. KRAFT AND C. PROCESI, On the geometry of conjugacy classes in classical groups, *Comment. Math. Helv.* **57** (1982), 539–602.
- [LS] T. LEVASSEUR AND S. P. SMITH, Primitive ideals and nilpotent orbits in type G_2 , *J. Algebra* **114** (1988), 81–105.
- [Ri] R. W. RICHARDSON, Derivatives of invariant polynomials on a semisimple Lie algebra, “Harmonic analysis and operator theory,” *Proc. Cent. Math. Anal. Aust. Natl. Univ.* **15** (1987), 228–241.
- [SK] M. SATO AND T. KIMURA, A classification of irreducible prehomogeneous vector spaces and their relative invariants, *Nagoya Math. J.* **65** (1977), 1–155.
- [Sh] T. SHOJI, On the Green polynomials of a Chevalley group of type F_4 , *Comm. Algebra* **10** (1982), 505–543.
- [Sl] P. SLODOWY, “Simple Singularities and Simple Algebraic Groups,” Lecture Notes in Mathematics, Vol. **815**, Springer-Verlag, Berlin/Heidelberg/New York, 1980.
- [VP] E. B. VINBERG AND V. L. POPOV, On a class of quasihomogeneous affine varieties, *Math. USSR-Izv.* **6** (1972), 743–758.